

ON n -GROUPOIDS

G. Čupona

An algebra $Q(f)$ with an n -ary operation is said to be an n -subgroupoid of a groupoid $G(\bullet)$ if $Q \subseteq G$ and f is the restriction of \bullet^{n-1} on Q . And, an algebra $A(F)$ is said to be an F -groupoid if there is a groupoid $G(\bullet)$ such that $A(f)$ is an n -subgroupoid of $G(\bullet)$ for every n -ary operator $f \in F$. It is shown in § 1 that every n -groupoid is an n -subgroupoid of a groupoid. The classes of n -subgroupoids of each of the classes of cancellative groupoids and commutative groupoids are described in §§ 2,3. It is shown in § 4 that the class of F -groupoids is a variety iff there is an n -ary operator $f \in F$ such that, for every m -ary operator $g \in F$, $n-1$ is a divisor of $m-1$.

1. Universal covering groupoids. An algebra $Q(f)$ with an n -ary operation is said to be an n -groupoid, and it is an n -subgroupoid of a groupoid $G(*)$ if $Q \subseteq G$ and

$$fa_1 \dots a_n = *^{n-1} a_1 \dots a_n,$$

for all $a_1, \dots, a_n \in Q$. The following result can be obtained as a corollary from the main results of the papers [4] and [6], but we shall give here a direct proof.

1.1. Every n -groupoid is an n -subgroupoid of a groupoid.

Proof. Let $Q(f)$ be an n -groupoid and $W(o)$ be the groupoid which is freely generated by the set Q . Thus, W is the minimal set of finite sequences on $Q \cup \{o\}$ (where $o \notin Q$) satisfying the following statements:

(i) $Q \subseteq W$; (ii) $u, v \in W \Rightarrow o uv \in W$.

Denote by U the set of elements of W in which do not occur subsequences of the following form:

$$o^{n-1} a_1 \dots a_n \quad (a_1, \dots, a_n \in Q).$$

It is easy to see that if $Q(f)$ is an n -subgroupoid of a cancellative (right cancellative) groupoid, then $Q(f)$ is cancellative (right cancellative) and the following quasiidentity is satisfied in $Q(f)$:

$$\begin{aligned} fx_1 \dots x_i z_1 \dots z_{n-i} = fy_1 \dots y_i z_1 \dots z_{n-i} &\Rightarrow \\ fx_1 \dots x_i u_1 \dots u_{n-i} = fy_1 \dots y_i u_1 \dots u_{n-i}, &\end{aligned} \quad (2.1)$$

for each $i \in \{1, \dots, n\}$.

Conversely, assume that $Q(f)$ is a right cancellative n -groupoid in which all the quasiidentities (2.1) are satisfied. The universal covering $U(\bullet)$ of $Q(f)$ can be not right cancellative. We are asking for a congruence α such that $Q(f)$ can be embedded as an n -subgroupoid in $U/\alpha(\bullet)$ and $U/\alpha(\bullet)$ should be right cancellative.

First, for each $i \in \{1, \dots, n-1\}$, let Q_i be defined by:

$$Q_i = \{o^{i-1} a_1 \dots a_i \mid a_1, \dots, a_i \in Q\},$$

and let α_i be a relation in Q_i defined by:

$$\begin{aligned} o^{i-1} a_1 \dots a_i \alpha_i o^{i-1} b_1 \dots b_i &\Leftrightarrow \\ (\exists c_{i+1}, \dots, c_n \in Q) fa_1 \dots a_i c_{i+1} \dots c_n = fb_1 \dots b_i c_{i+1} \dots c_n. &\end{aligned}$$

By (2.1), the quantifier \exists may be changed by \forall , and this implies that:

$$c \in Q, o^{i-1} a_1 \dots a_i \alpha_i o^{i-1} b_1 \dots b_i \Rightarrow o^i a_1 \dots a_i c \alpha_{i+1} o^i b_1 \dots b_i c.$$

We also note that α_1 is the equality on $Q (= Q_1)$.

Denote by α the minimal relation on U which satisfy the following propositions:

$$\alpha_1 \cup \alpha_2 \cup \dots \cup \alpha_{n-1} \subseteq \alpha,$$

$$u_1 \alpha v_1, u_2 \alpha v_2, o u_1 u_2, o v_1 v_2 \in U \Rightarrow o u_1 u_2 \alpha o v_1 v_2.$$

It is easy to see that α is a congruence on $U(\bullet)$ and that $G(\bullet) = U/\alpha(\bullet)$ is a right cancellative groupoid. Moreover, if $Q(f)$ is cancellative, then $G(\bullet)$ is cancellative too.

Finally, the mapping $a \rightarrow a^\alpha$ embeds $Q(f)$ into $G(\bullet)$.

Thus we obtain the following result.

2.2. *An n -groupoid $Q(f)$ is an n -subgroupoid of a (right) cancellative groupoid iff $Q(f)$ is (right) cancellative and satisfies all the quasiidentities (2.1). \square*

As a corollary of 2.2 and the fact that every groupoid with cancellation is a subgroupoid of a quasigroup ([1], VII. 4) we obtain the following result.

2.3. *The class of n -subgroupoids of quasigroups and the class of n -subgroupoids of groupoids with cancellation are equal. \square*

If $n \geq 3$, then there exist n -quasigroups which do not satisfy some of the quasiidentities (2.1) (for example, [7] p. 115), and thus we get the following result.

2.4. If $n \geq 3$, then there exist n -quasigroups which are not n -subgroupoids of quasigroups. \square

3. Commutative n -groupoids. An n -groupoid $Q(f)$ is said to be (i, j) -commutative if:

$$fx_1 \dots x_i \dots x_j \dots x_n = fx_1 \dots x_j \dots x_i \dots x_n$$

is an identity equation. $Q(f)$ is called commutative if it is (i, j) -commutative for each pair (i, j) : $1 \leq i < j \leq n$.

3.1. *An n -groupoid $Q(f)$ is an n -subgroupoid of a commutative groupoid iff $Q(f)$ is $(1, 2)$ -commutative.*

Proof. Let $Q(f)$ be a $(1, 2)$ -commutative n -groupoid, and let $C(o)$ be the freely generated commutative groupoid by the set Q . Then

$$o uv = o u' v' \Leftrightarrow (u = u', v = v') \text{ or } (u = v', v = u').$$

Denote by D the set of elements of C which can not be represented as products of the form:

$$\Pi (a_1, \dots, a_{t-1}, o^{n-1} b_1 \dots b_n, a_{t+1}, \dots, a_m)$$

with $a_t, b_j \in Q$.

Define a binary operation \bullet on D by:

$$u, v, o uv \in D \Rightarrow \bullet uv = o uv,$$

and

$$u, v \in D, o uv = o^{n-1} a_1 \dots a_n, a = fa_1 \dots a_n \Rightarrow \bullet uv = a.$$

It is easy to see that:

- (i) the operation \bullet is well defined;
- (ii) $D(\bullet)$ is a commutative groupoid;
- (iii) $Q(f)$ is an n -subgroupoid of $D(\bullet)$.

It is clear that every n -subgroupoid of a commutative groupoid is a (1,2)-commutative n -groupoid. \square

A groupoid $G(*)$ is said to be n -commutative if the n -groupoid $G(*^{n-1})$ is commutative.

3.2. *The class of n -subgroupoids of n -commutative groupoids and the class of commutative n -groupoids are equal.*

Proof. First, it is clear that every n -subgroupoid of an n -commutative groupoid is a commutative n -groupoid.

Let $Q(f)$ be a commutative n -groupoid and let $U(\bullet)$ be the universal covering groupoid of $Q(f)$. Define a relation α on U in the following way. If $v \rightarrow i_v$ is a permutation of $\{1, \dots, n\}$, Π a product on $U(\bullet)$, and $u_i, t_j \in U$, then:

$$\Pi (u_1, \dots, u_{p-1}, \bullet^{n-1} t_1 \dots t_n, u_{p+1}, \dots) \alpha$$

$$\Pi (u_1, \dots, u_{p-1}, \bullet^{n-1} t_{i_1} \dots t_{i_n}, u_{p+1}, \dots).$$

It is obvious that the transitive and reflexive extension β of the relation α is a congruence on $U(\bullet)$ and that the groupoid $U/\beta(\bullet) = G(\bullet)$ is n -commutative.

From **1.2** and **1.3** it follows that:

$$a \in Q, u \in U \Rightarrow (a \alpha u \Rightarrow a = u),$$

and this implies that:

$$a, b \in Q \Rightarrow (a \beta b \Rightarrow a = b),$$

i.e. that $Q(f)$ can be embedded in $G(\bullet)$ as an n -subgroupoid. \square

The statements **3.1** and **3.2** imply that every commutative n -groupoid is an n -subgroupoid of a commutative groupoid, and also an n -subgroupoid of an n -commutative groupoid. But there exist commutative n -groupoids which can not be embedded in groupoids which are both commutative and n -commutative, for commutativity and n -commutativity imply some associativity. (For example, every commutative and 3-commutative groupoid is a semigroup.)

4. F-groupoids. Here we assume that F is a nonempty set of finitary operators such that $F_0 \cup F_1 = \emptyset$, where F_n is the set of n -ary operators belonging to F . An algebra $A(F)$ is said to be an F -groupoid if there is a groupoid $G(*)$ such that $A(f)$ is an n -subgroupoid of $G(*)$ for every n -ary operator $f \in F$; then we also say that $A(F)$ is an F -subgroupoid of $G(*)$. An algebra $A(F)$ is said to be a weak F -groupoid if for every sequence of operators $f_1, \dots, f_r, g_1, \dots, g_s \in F$ such that:

$$f_i \in F_{n_i+1}, g_j \in F_{m_j+1}, n_1 + \dots + n_r = n = m_1 + \dots + m_s \quad (4.1)$$

the following identity is satisfied in $A(F)$:

$$f_1 \dots f_r x_0 \dots x_n = g_1 \dots g_s x_0 \dots x_n. \quad (4.2)$$

4.1. Every F -groupoid is a weak F -groupoid.

Proof. Let $A(F)$ be an F -subgroupoid of a groupoid $G(*)$, and assume that (4.1) is satisfied. If $a_0, \dots, a_n \in A$ then we have:

$$f_1 \dots f_r a_0 \dots a_n = *^n a_0 \dots a_n = g_1 \dots g_s a_0 \dots a_n,$$

i.e. (4.2) is an identity in $A(F)$. \square

Let J_F be the following set of integers:

$$J_F = \{n \mid F_{n+1} \neq \emptyset\},$$

and denote by d_F the greatest common divisor of the numbers belonging to J_F .

4.2. Every weak F -groupoid is an F -groupoid iff $d_F \in J_F$.

Proof. Let $d = d_F \in J_F$, $f \in F_{d+1}$ and let $A(F)$ be a weak F -groupoid. By 1.1, $A(f)$ is a $d+1$ -subgroupoid of a groupoid $U(\bullet)$. If $g \in F_{m+1}$, then d is a divisor of m , and by (4.2) we have:

$$g x_0 \dots x_m = f^{m/d} x_0 \dots x_m = \bullet^m x_0 \dots x_m,$$

and this implies that $A(F)$ is an F -subgroupoid of $U(\bullet)$.

Assume now that $d_F \notin J_F$. Then, if n is the least element of J_F there is an element $m \in J_F$ which is not divisible by n , and we shall assume that m is the least element of J_F with that property. Define an algebra $A(F)$ in the following way:

- (i) $A = \{a, b, c\}$, $a \neq b \neq c \neq a$;
- (ii) $f \in F_{m+1} \Rightarrow f a_0 \dots a_m = a$ if $a_v \neq c$ for some v and $f c^{m+1} = b$;
- (iii) $g \in F_{k+1}$, $k \neq m \Rightarrow g a_0 \dots a_k = a$.

It is easy to see that $A(F)$ is a weak F -groupoid. $A(F)$ is not an F -groupoid, for if $A(F)$ were an F -subgroupoid of a groupoid $G(*)$ and if $f \in F_{m+1}$, $g \in F_{n+1}$ then we would have:

$$\begin{aligned} b &= f c^{m+1} = *^{m-n} g c^{m+1} = *^{m-n} a c^{m-n} = \\ &= *^{m-n} g a^{n+1} c^{m-n} = *^m a^{n+1} c^{m-n} = f a^{n+1} c^{m-n} \\ &= a. \quad \square \end{aligned}$$

If Σ is a class of groupoids we can ask for an axiom system of the class of F -subgroupoids of Σ -groupoids. We note that there are known convenient descriptions of F -subgroupoids of semigroups ([2], 5), and F -subgroupoids of cancellative semigroups ([5], 3).

REFERENCES

- [1] P. M. Cohn, Universal algebra, Harper & Row, New York 1965.
- [2] N. Celakoski, On semigroup associatives, Maced. Acad. of Sc. and Arts; Contributions IX — 2 (1977), 5—19.
- [3] G. Čupona, S. Markovski, On quasigroups, Godišen zbornik Prir.-matem. fakultet Skopje, 25/26 (1975/76) Sekc. A, 9—14.
- [4] M. D. Prešič, S. B. Prešič, Od the embedding of Ω -algebras in groupoids, Publications de l'Institut Mathématique N. S. T. 21 (35), 1977, 169—174.
- [5] Ā. Чупона, Ассоцијативи со кратење, Годишен зборник Прир.-матем. факултет Скопје, 19 (1969) Секц. А, 5—14.
- [6] Ā. Чупона, С. Марковски, Сместување на универзални алгебри, Годишен зборник Прир.-матем. факултет Скопје, 25/26 (1975/76) Секц. А, 15—34.
- [7] В. Д. Белоусов, n -арные квазигруппы, Кишинев 1972.

Ā. Чупона

ЗА n -ГРУПОИДИТЕ

(Резиме)

Во работава се покажува дека секој n -групоид е n -подгрупоид на групоид. Се дава опис на класата n -подгрупоиди на групоиди со кратење како и на класата n -подгрупоиди од комутативни групоиди. Се разгледува и поопштото прашање за сместување на произволни алгебри во групоиди и се докажува дека класата F -групоиди е многократност акко постои n -арен оператор $f \in F$ таков што $n - 1$ е делител на $m - 1$ за секој m -арен оператор $g \in F$.