

n-SUBSEMIGROUPS OF SOME COMMUTATIVE SEMIGROUPS

G. Čupona

A subset Q of a semigroup S is said to be an n -subsemigroup of S if: $a_1, \dots, a_n \in Q \Rightarrow a_1 \dots a_n \in Q$. Then, by $[x_1 \dots x_n] = x_1 \dots x_n$ is defined an associative n -ary operation $[]$ on Q , i.e. $(Q; [])$ is an n -semigroup. If C is a class of semigroups, then the class of n -semigroups that are n -subsemigroups of C -semigroups is denoted by $C(n)$. A variety of semigroups C is called an n -variety of semigroups iff $C(n)$ is a variety of n -semigroups. (Clearly, every variety of semigroups is a 2-variety.) The set of n -varieties of semigroups and its complement, in the set of varieties of semigroups, are infinite for any $n > 3$. ([1], [2], [3], [4]) It is also known that the set of n -varieties of commutative semigroups is infinite. ([2]) Here we show that if $n > 3$ then the set of varieties of commutative semigroups that are not n -varieties is also infinite.

Further on, by a semigroup (n -semigroup) we will mean a commutative semigroup (commutative n -semigroup).

A variety of semigroups is defined by a set of identities of the following form:

$$x_1^{i_1} x_2^{j_2} \dots x_p^{j_p} = x_1^{j_1} x_2^{i_2} \dots x_p^{i_p}, \quad (*)$$

where: x_1, x_2, \dots are variables; i_v, j_v are nonnegative integers such that $(\sum i_v)(\sum j_v) > 0$. If $i_v = j_v$, for each v , then $(*)$ is called a trivial identity. (As usually, variables will be also denoted by x, y, z, \dots).

Let m, s, n, k be positive integers such that $s + 2 \leq m, m \neq 2s + 1, m \neq 2s + 2, m \equiv 1 \pmod{n-1}$ and $m + 2 \leq k$. Consider the following two identities:

$$x^s y^{m-s} = x^{s+2} y^{m-s-1}, \quad (m, s)$$

$$x_1 x_2 \dots x_k = x_1^2 x_2 \dots x_k. \quad (k)$$

Denote by $(m, s; k)$ the set of the given two identities, and by $(m, s; k)^n$ the set of identities $(*)$ which are consequences of $(m, s; k)$ and the exponents satisfy the following condition:

$$\sum i_v \equiv \sum j_v \equiv 1 \pmod{n-1}.$$

Every identity (*) which belongs to $(m, s; k)^n$ induces an identity

$$[x_1^{i_1} \cdots x_p^{i_p}] = [x_1^{j_1} \cdots x_p^{j_p}] \quad [*]$$

of n -semigroups. We denote by $[m, s; k]^n$ the set of identities [*] such that (*) is in $(m, s; k)^n$.

It is clear that if $x^s y^{m-s} = x^i y^j$ is in $(m, s; k)^n$, then $i = s, j = m - s$, i.e. $[m, s; k]^n$ does not contain a nontrivial identity $[x^s y^{m-s}] = [x^i y^j]$.

Consider the variety $C^{(m, s; k)}$ of semigroups defined by $(m, s; k)$. We will show namely that $C^{(m, s; k)}$ is not an n -variety. To prove this statement it is enough to find an n -semigroup $(Q; [\])$ which satisfies all identities belonging to $[m, s; k]^n$, but does not belong to $C^{(m, s; k)}(n)$.

Let a, b, c be three different objects and let $(Q; [\])$ be the n -semigroup with a presentation

$$\langle a, b, c; [a^{s+2} c^{m-s-2}] = [b^{s+2} c^{m-s-2}] \rangle \quad (**)$$

in the variety of n -semigroups defined by $[m, s; k]^n$. Let us give a more explicit construction of $(Q; [\])$. First, let $(F; [\])$ be the free n -semigroup with a basis $B = \{a, b, c\}$ in the variety defined by $[m, s; k]^n$. In other words F consists of all „commutative products of powers“ $[a^i b^j c^p]$, such that: $i, j, p \geq 0, i + j + p \equiv 1 \pmod{n-1}$, and the equality

$$[a^j b^j c^p] = [a^{j'} b^{j'} c^{p'}]$$

holds in F iff the following identity

$$[x^i y^j z^p] = [x^{i'} y^{j'} z^{p'}]$$

is in $[m, s; k]^n$. The operation $[\]$ is defined in the usual way, i.e. by the following equation:

$$[[a^{i_1} b^{j_1} c^{p_1}] \cdots [a^{i_n} b^{j_n} c^{p_n}]] = [a^{i_1 + \cdots + i_n} b^{j_1 + \cdots + j_n} c^{p_1 + \cdots + p_n}].$$

Consider the minimal congruence \approx on $(F; [\])$ such that

$$[a^{s+2} c^{m-s-2}] \approx [b^{s+2} c^{m-s-2}].$$

Namely \approx is defined in the following way. If $u = a^i b^j c^k$ is such that $i + j + k \equiv 0 \pmod{n-1}$, then:

$$[u a^{s+2} c^{m-s-2}] \sim [u b^{s+2} c^{m-s-2}] \text{ and } [u b^{s+2} c^{m-s-2}] \sim [u a^{s+2} c^{m-s-2}].$$

Now, \approx is the transitive and reflexive extension of \sim , i.e. $u \approx v$ iff there exist $u_0, \dots, u_t \in F$ such that $u = u_0, v = u_t, t \geq 0$, and $u_{i-1} \sim u_i$ if $i \geq 1$. Then $(F/\approx; [\])$ is the desired n -semigroup $(Q; [\])$. We can assume that $a, b, c \in Q$.

Let us show that $[a^s c^{m-s}] \neq [b^s c^{m-s}]$ in $(Q; [])$. Namely, we first conclude that if $[a^s c^{m-s}] = [a^i b^j c^p]$ in F , then $i = s, j = 0, p = m - s$. We also have that $[a^s c^{m-s}] \sim' [a^i b^j c^p]$, for any i, j, p , and therefore $[a^s c^{m-s}] \approx' [b^s c^{m-s}]$, i.e. $[a^s c^{m-s}] \neq [b^s c^{m-s}]$ in $(Q; [])$.

Now, it is easy to show that $(Q; [])$ does not belong to $C^{(m, s; k)}(n)$. Namely, if $(Q; [])$ were an n -subsemigroup of a semigroup $S \in C^{(n, s; k)}$, then we would have

$$\begin{aligned} a^s c^{m-s} &= a^{s+2} c^{m-s-1} \\ &= a^{s+2} c^{m-s-2} c \\ &= b^{s+2} c^{m-s-2} c \\ &= b^{s+2} c^{m-s-1} \\ &= b^s c^{m-s} \end{aligned}$$

in S , and this would imply the equality $[a^s c^{m-s}] = [b^s c^{m-s}]$ in $(Q; [])$.

Clearly, if $m + 2 \leq k' < k''$ then $C^{(m, s; k')}$ is a proper subvariety of $C^{(m, s; k'')}$, and thus if s and m are fixed positive integers such that $s + 2 \leq m$, $m \not\equiv 2s + 1$, $m \not\equiv 2s + 2$, $m \equiv 1 \pmod{n-1}$ then we have an infinite set of varieties $\{C^{(m, s; k)} \mid k \geq n + 2\}$ of commutative semigroups which are not n -varieties.

Denote by $C^{(m, s)}(C^{(k)})$ the variety of semigroups defined by the identity $(m, s) ((k))$. From the above considerations it follows that $C^{(m, s)}$ is not an n -variety. Namely, we notice again that there is not a nontrivial identity $[x^s y^{m-s}] = [x^i y^j]$ in $[m, s]^n$. And, the n -semigroup $(Q; [])$ with a presentation $(**)$ in the variety of n -semigroups defined by $[m, s]^n$ is not an n -subsemigroup of a semigroup belonging to $C^{(m, s)}$.

But, the variety $C^{(k)}$ is an n -variety for every pair of positive numbers n, k such that $n \geq 2$. (Namely, the assumption $k \geq n + 2$ is not necessary.) First we notice that $(*)$ is a consequence of (k) if $i_v = j_v$ for each v or $\sum i_v \geq k$, $\sum j_v \geq k$ and $i_v > 0 \Leftrightarrow j_v > 0$. This gives a complete description of $[k]^n$, as well.

Assume now that $(Q; [])$ is an n -semigroup which satisfy any identity of $[k]^n$, i.e. each identity $[*]$ such that

$$\sum j_v \equiv \sum j_v \equiv 1 \pmod{n-1}, \quad \sum i_v \geq k, \quad \sum j_v \geq k$$

and $i_v > 0 \Leftrightarrow j_v > 0$.

We have to show that $(Q; [])$ is an n -subsemigroup of a semigroup belonging to $C^{(k)}$.

Let $n \geq k$, and let a binary operation \cdot be defined on Q by:

$$x \cdot y = [x y^{n-1}].$$

Then it is easy to see that $(Q; \cdot)$ is a semigroup in $C^{(k)}$, and moreover that

$$x_1 x_2 \dots x_n = [x_1 x_2 \dots x_n],$$

and therefore $(Q; [\])$ is an n -subsemigroup of $(Q; \cdot)$.

In the general case, we consider the semigroup S with a presentation

$$\langle Q; \{a = a_1 \dots a_n \mid a = [a_1 \dots a_n] \text{ in } (Q; [\])\} \rangle$$

in the variety $C^{(k)}$, and we have to show that:

$$a, b \in Q \Rightarrow (a = b \text{ in } S \Rightarrow a = b \text{ in } Q),$$

but here we will not give the complete proof of this statement.

REFERENCES

- [1] Марковски С.: За дистрибутивните полугрупи, Год. збор. Матем. фак. Скопје 30 (1979), 15—27
- [2] Марковски С.: За една класа полугрупи, Математички Билтен Кн. 1 (XXVII) 29—36
- [3] Чупона Г.: Полугрупи генерирани од асоцијативи. Год. зборн. ПМФ — Скопје Секц. А 15 (1964), 5—25
- [4] Чупона Г.: n -subsemigroups of semigroups satisfying the law $x^r = x^{r+m}$, Год. зборник Матем. фак. Скопје, 30 (1979), 5—14

Г. Чупона

n — ПОТПОЛУГРУПИ ОД НЕКОИ КОМУТАТИВНИ ПОЛУГРУПИ

Резиме

Во работава се покажува дека множеството многубразија M комутативни полугрупи, такви што $M(n)$ (т.е. класата од n -потполугрупи од M -полугрупи) е вистинско квазимногубразие, е бесконечно, за секое $n > 3$.